

Anti-IL-4 [11B11] Vivopure 10 mg, 10 mg, Ab00147-6.4-VPT View online

Anti-IL-4 [11B11] Vivopure 10 mg Ab00147-6.4-VPT

This antibody was created using our proprietary Fc Silent[™] engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

Isotype and Format: Rat IgG1, Fc Silent[™], Kappa

Clone Number: 11B11

Alternative Name(s) of Target: Interleukin-4; II4; II-4; B-cell IgG differentiation factor; B-cell growth factor 1; B-cell stimulatory factor 1; BSF-1

UniProt Accession Number of Target Protein: P07750

Published Application(s): in vivo, ELISA, FC, IHC

Published Species Reactivity: Mouse

Immunogen: BSF-1 / IL-4 purified from serum-free supernatants of cultured EL-4 cells (a C57BL/6 thymoma) induced with 4 β -phorbol 12 β -myristate 12 α -acetate for 48h.

Specificity: Binds to a core functional region of murine IL-4, which prevents this cytokine from interacting with its cognate receptor (Yao et al., 2006 PMID: 16361318).

Application Notes:

Antibody First Published in: Ohara J, Paul WE. Production of a monoclonal antibody to and molecular characterization of B-cell stimulatory factor-1. Nature. 1985 May 23-29;315(6017):333-6. PMID:2582266 **Note on publication:** Describes the isolation of the monclonal antibody and its use in studying IL-4 (before it was named IL-4).

Product Form

Size: 10 mg Vivopure products are produced at high purity (>98%), low endotoxin (<0.5 EU/mg) and are formulated without preservatives. As a result Vivopure products are the ideal choice for in vivo research applications.

Purification: Protein A affinity purified

Supplied In: PBS only, with >98% antibody purity and <1 EU/mg guaranteed.

Storage Recommendation: All vivopure products are formulated in PBS only without addition of preservatives. To ensure optimal storage and prevent microbial contamination, only open and dispense under sterile conditions.

Concentration: >=1mg (see vial label for exact conc)

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.