

## Anti-Rhodopsin [Rho 1D4] Bulk Size Ab00337-1.4-BT

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

**Isotype and Format:** Mouse IgG1, Fc Silent<sup>™</sup>, Kappa

Clone Number: Rho 1D4

Alternative Name(s) of Target: Rho-tag; Visual Purple UniProt Accession Number of Target Protein: P08100

Published Application(s): IP, Radioimmune labeling, WB, ELISA, IF, IHC

Published Species Reactivity: Cow, Human, Zebrafish, Amphibians, Rat, Mouse

**Immunogen:** Bleached bovine rod outer segment (ROS).

**Specificity:** The antibody binds to the C-terminus of rhodopsin on the cytoplasmic side of disk membranes. The epitope has the following amino acid sequence: 'TETSQVAPA'. A limitation of the 1D4 tag, however, is that it has to be placed at the C-terminus of a protein. This is because the Rho1D4 monoclonal antibody requires a free carboxylate group for high affinity binding. Amidation of the carboxyl group lowers its immunoreactivity to the Rho1D4 antibody by over 100 fold.

**Application Notes:** The antibody binds to the C-terminus of rhodopsin on the cytoplasmic side of disk membranes. Rhodopsin is the major pigment in rod photoreceptors and belongs to the family of G-protein-coupled receptors (GPCRs). The linear epitope (TETSQVAPA) can be used as a protein tag.

**Antibody First Published in:** Molday et al. 1983 Monoclonal antibodies to rhodopsin: characterization, corss-reactivity, and application as sturctural probes. Biochemistry, 1983; 22(3):653-660 PMID:6188482 **Note on publication:** Describes the generation of two monoclonal antibodies from hybridoma.

## **Product Form**

**Size:** 1 mg Purified antibody in bulk size. **Purification:** Protein A affinity purified

Supplied In: PBS only.

**Storage Recommendation:** Store at 4°C for up to 3 months. Note, this antibody is provided without added preservatives, it is therefore recommed this antibody be handled under sterile conditions. For longer

storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for r procedures for humans or animals.	research use only.	It is not intended fo	or use in therapeutic	or diagnostic