

## Anti-Botulinum neurotoxin [MCS6-27] Standard Size Ab00508-1.1

**Isotype and Format:** Mouse IgG1, Kappa

**Clone Number:** MCS6-27

**Alternative Name(s) of Target:** BBoNT/B

**UniProt Accession Number of Target Protein:**

**Published Application(s):** WB, ELISA

**Published Species Reactivity:** Clostridium botulinum

**Immunogen:** Recombinant BoNT/B fragments expressed in E. coli as GST-fusion proteins.

**Specificity:**

**Application Notes:** The antibody binds specifically to BoNT/B, a two-chain neurotoxic protein produced by the bacterium Clostridium botulinum. Botulism, an infection of the bacterium, which is commonly spread through contaminated food, can be a fatal disease, and symptoms include a gradual flaccid paralysis. Paralysis is caused by the internalization by endocytosis of the toxin in specific axons, mediated by the attachment of the heavy chain of the protein to axon surface. Once inside the axon, BoNT/B acts by cleaving SNARE proteins, such as VAMP, which are involved in the docking and fusion of neurosecretory vesicles. This prevents the release of the neurotransmitter acetylcholine, resulting in paralysis of the muscles. The monoclonal antibody MCS6-27 binds to the Hc and H5 fragments of BoNT/B, and the binding epitope is located in the heavy chain, between amino acids E859 and E1291. The monoclonal antibody is able to capture BoNT/B at SDS concentrations between 0 and 0.4 mM, and can be used to screen food samples for botulinum toxin.

**Antibody First Published in:** Scotcher MC, Cheng LW, Stanker LH. Detection of botulinum neurotoxin serotype B at sub mouse LD(50) levels by a sandwich immunoassay and its application to toxin detection in milk. PLoS One. 2010 Jun 10;5(6). [PMID:20548779](#)

**Note on publication:** Describes the characterization of monoclonal antibodies against BoNT/B, and the generation of a sensitive sandwich immunoassay.

## Product Form

**Size:** 200 µg Purified antibody.

**Purification:** Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

**Concentration:** 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.