

Anti-CD44 [OX-49] Standard Size Ab00550-2.3

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

This reformatted mouse antibody was made using the variable domain sequences of the original Mouse IgG2a format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Mouse IgG2a, [Fc Silent™](#), Kappa

Clone Number: OX-49

Alternative Name(s) of Target: CDW44, CSPG8, ECMR-III, HCELL, HUTCH-I, IN, LHR, MC56; MDU2; MDU3; MIC4; Pgp1; CD44 molecule (Indian blood group); HCAM; Hermes antigen; Pgp-1; lymphocyte homing receptor; ECM-III; HUTCH-1

UniProt Accession Number of Target Protein:

Published Application(s): IP, WB, FC

Published Species Reactivity: Rat

Immunogen: T blasts from a mixed lymphocyte reaction between purified rat CD4+ T cells and irradiated spleen.

Specificity: This antibody binds rat CD44 at an epitope overlapping with asparagine residue 67. The murine homologue of CD44 has a histidine in this position which is sufficient to abrogate binding (Zheng et al. 1995).

Application Notes: This antibody antibody recognises rat CD44.

Antibody First Published in: Paterson et al. Antigens of activated rat T lymphocytes including a molecule of 50,000 Mr detected only on CD4 positive T blasts. Mol Immunol. 1987 Dec;24(12):1281-90. [PMID:2828930](#)

Note on publication: Describes the generation and initial characterisation of this antibody.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration:

1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.