

Anti-BORIS [20B11] Standard Size Ab00825-5.0

Developed in partnership with Ximbio (www.ximbio.com).

Isotype and Format: Mouse IgG3, Kappa

Clone Number: 20B11

Alternative Name(s) of Target: CT27; Brother of the regulator of imprinted sites; Cancer/testis antigen 27; CCCTC binding factor (zinc finger protein) like; CCCTC-binding factor; CTCF paralog; CTCF T; CTCF-like protein; Ctcfl; CTCFL_HUMAN; dj579F20.2; HMG 1L1; HMGB1L1; MGC163358; MGC169105; MGC169106; CTCFL

UniProt Accession Number of Target Protein: Q8NI51

Published Application(s): CHIP, IP, WB, ELISA, IF, IHC

Published Species Reactivity: Human

Immunogen: Synthetic peptide within the BORIS C-terminal domain (aa 614-648) (CG)GEMFPVACRETTARVKEE (NB - the first two aa do not belong to BORIS)

Specificity: 20B11 reacts specifically with BORIS.

Application Notes: Reagent for research, diagnostic tool. BORIS protein has been identified as Cancer-Testis Antigen (CTA) with testis-specific paralogue of the CCCTC-binding factor. Recent studies have demonstrated that d BORIS is directly responsible for the transcriptional activation of TSP50 (testes-specific protease 50)

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.