

## Anti-CD27 [RM27-3E5] VivopureX 5 mg Ab00895-1.1-VXM

This chimeric mouse antibody was made using the variable domain sequences of the original Rat IgG2a format, for improved compatibility with existing reagents, assays and techniques.

**Isotype and Format:** Mouse IgG1, Kappa

**Clone Number:** RM27-3E5

**Alternative Name(s) of Target:** TNFRS7; CD27 antigen; CD27L receptor; T-cell activation antigen CD27; Tumor necrosis factor receptor superfamily member 7

**UniProt Accession Number of Target Protein:** P41272

**Published Application(s):** IP, Neutralisation, FC

**Published Species Reactivity:** Mouse

**Immunogen:** This antibody was raised by immunising mouse with mouse CD27-human IgG1 Fc fusion protein. Isolated popliteal lymph node cells were then fused with P3U1 myeloma cells to produce stable hybridomas.

**Specificity:** This antibody is specific for the extracellular domain of murine CD27, a TNF-receptor superfamily member.

**Application Notes:** This antibody recognises murine CD27, as confirmed by flow-cytometry and immunoprecipitation analysis (Sakanishi & Yagita, 2010), a member of the TNF-receptor superfamily. The interaction between CD27 and its ligand CD70, which is recognised by anti-CD70 antibody TAN 1-7 (Ab00816), is crucial to regulation of the survival and differentiation of T cells, B cells and NK cells. As a result, RM27-3E5 modulates multiple immune cell types. The original rat IgG2a version of RM27-3E5 did not deplete T cells, NKT or NK cells, but does enhance the proliferation of anti-CD3-stimulated splenic T cells (Sakanishi & Yagita, 2010). In murine models of melanoma, it promotes the persistence of tumour-specific CD8+ T cells within tumours, and reduces levels of PD-1 expression on CD8+ T cells (Roberts et al, 2010). Additionally, anti-CD27 reduces the proportion of FoxP3+ CD4+ T cells in tumours, and enhances the effector activity of IFN $\gamma$ -secreting, tumour-infiltrating CD8+ T cells and NK cells (Roberts et al, 2010). Finally, anti-CD27 Ab treatment alters murine splenocyte cytokine production, significantly reducing IL-5 and IFN- $\gamma$  synthesis, and increasing TNF- $\alpha$  production (Sumi et al, 2008). It also significantly suppresses total serum IgE levels, while significantly increasing IgG1 and IgG2a levels. These immune modulatory properties translate into potent anti-tumour effects in mice. Treatment of established murine melanoma with RM27-3E5 (rat IgG2a) resulted in a significant reduction in lung metastases and subcutaneous tumours, through an IFN $\gamma$ -dependent mechanism mediated by CD8+ T cells and NK cells (Roberts et al,

2010). In murine models of T cell lymphoma, the tumour-specific CTL response induced by RM27-3E5 (rat IgG2a) promoted almost complete regression of tumours, regardless of CD27 expression on tumour cells (Sakanishi & Yagita, 2010). This antibody has been used to identify CD27-expressing cells through flow-cytometry (Koyanagi et al, 2012).

**Antibody First Published in:** Sumi et al CD27 and CD70 do not play a critical role in the development of experimental allergic conjunctivitis in mice Immunology Letters, Volume 119, Issues 1-2, 15 August 2008, Pages 91-96 [PMID:18579220](https://pubmed.ncbi.nlm.nih.gov/18579220/)

**Note on publication:** Describes the original generation of this antibody.

## Product Form

**Size:** 5 mg VivopureX products are produced at high purity (>98%), low endotoxin (<0.5 EU/mg) and are formulated without preservatives. These antibodies are chimerized to have an Fc domain matching their target species to reduce immunogenicity and give you the optimal effector function for your experiment. As a result VivopureX products are the ideal choice for in vivo research applications.

**Purification:** Protein A affinity purified

**Supplied In:** PBS only.

**Storage Recommendation:** All VivopureX products are formulated in PBS only without addition of preservatives. To ensure optimal storage and prevent microbial contamination, only open and dispense under sterile conditions. Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

**Concentration:** >=1mg (see vial label for exact conc)

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.