

Anti-CD74 [LN-2] Standard Size, 200 µg, Ab01546-13.12 View online

## Anti-CD74 [LN-2] Standard Size Ab01546-13.12

This chimeric human antibody was made using the variable domain sequences of the original Mouse IgG1 format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Human IgG4-S228P, Kappa

Clone Number: LN-2

Alternative Name(s) of Target: la antigen; HLA class II histocompatibility antigen gamma chain; HLA-DR antigens-associated invariant chain; la antigen-associated invariant chain; li; p33; LN2

UniProt Accession Number of Target Protein: P04233

Published Application(s): IP, WB, ELISA, FC, IF, IHC

Published Species Reactivity: Rat, Human, Mouse

Immunogen: This antibody was raised by immunizing mice with SU-DHL-4 lymphoma cells.

**Specificity:** LN-2 antibody recognizes CD74. It reacts with an epitope residing within 60 amino acids of the extracytoplasmic, COOH terminus of the protein. CD74 is a type II transmembrane protein which binds to the peptide binding groove of newly synthesized MHC class II alpha/beta heterodimers and prevents their premature association with endogenous polypeptides.

**Application Notes:** Anti-CD74 has been shown to be useful in differentiating atypical fibroxanthoma (-) from malignant fibrous histiocytoma (+). This antibody can be used for the identification of B cell lymphoma and leukemias. It also stains activated neoplastic cells in T cell lymphomas and Sternberg-reed cells. Furthermore, this antibody was shown to be capable of distinguishing small cell lung cancer from a non small cell lung cancer as a part of a panel of antibodies (loachim et al., 1996). Furthermore, LN2 was successfully used in immunohistochemical and flow cytometric analysis of lung cancer tissue and cells (loachim et al., 1996). Finally, this antibody is also useful in immunofluorescence analysis of CD74, as it was very effective in labelling this protein in CD74-transfected COS-7 cells (Leng et al., 2003).

**Antibody First Published in:** Epstein et al. Two new monoclonal antibodies (LN-1, LN-2) reactive in B5 formalin-fixed, paraffin-embedded tissues with follicular center and mantle zone human B lymphocytes and derived tumors. J Immunol. 1984 Aug;133(2):1028-36. PMID:6376628

**Note on publication:** This article describes the LN-2 antibody.

## **Product Form**

Size: 200 µg Purified antibody. Purification:

© 2024 Absolute Antibody Wilton, UK.

Protein A affinity purified

**Supplied In:** PBS with 0.02% Proclin 300.

**Storage Recommendation:** Store at 4°C for up to 3 months. For longer storage, aliquot and store at - 20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.