

Anti-Spike Protein [CV1] Standard Size Ab02018-10.3

This antibody was created using our proprietary Fc Silent™ engineered Fc domain containing key point mutations that abrogate binding to Fc gamma receptors.

Isotype and Format: Human IgG1, [Fc Silent™](#), Lambda

Clone Number: CV1

Alternative Name(s) of Target: S-protein; SARS-CoV2; SARS Coronavirus 2; SARS-CoV-2; SARS CoV 2; 2019-nCoV; Severe acute respiratory syndrome coronavirus 2; Spike protein; S protein; SARS-CoV 2 S protein; S glycoprotein; E2; Peplomer protein; SARS-CoV2

UniProt Accession Number of Target Protein: P0DTC2

Published Application(s): neutralize, therapeutic, ELISA, FC

Published Species Reactivity: SARS Coronavirus 2

Immunogen: The original antibody was generated by isolating B-cells from the peripheral blood mononuclear cells (PBMCs) of SARS CoV2 infected patients, after the serum neutralizing activity was confirmed.

Specificity: This antibody binds the ectodomain encoded by amino acid residues 1-1208 of the SARS CoV 2 spike protein.

Application Notes: This antibody is recommended for diagnosis and/or treatment of SARS CoV 2 or 2019-nCoV. CV1 binds an epitope outside the SARS CoV 2 RBD was found to be weakly neutralizing (IC50=15µg/ml). ELISA was used to confirm the binding of SARS CoV 2 positive serum samples to the entire ectodomain and the receptor binding domain of the spike protein. Specific binding of the mAb to SARS CoV 2 to ectodomain was confirmed with biolayer interferometry. Phycoerythrin labelled mAb was used to stain 293E cells transfected with wild type SARS CoV 2 and were analysed by flow cytometry (Seydoux et al., 2020).

Antibody First Published in: Seydoux et al. Characterization of neutralizing antibodies from a SARS-CoV-2 infected individual [PMID:32511342](#)

Note on publication: Describes the generation and characterization of this antibody from PBMCs of infected patients.

Product Form

Size: 200 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In:

PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.