

Anti-Spike protein [MR3] Standard Size Ab02059-10.159

This chimeric human antibody was made using the variable domain sequences of the original VHH format, for improved compatibility with existing reagents, assays and techniques.

Isotype and Format: Human IgG1-Fc fusion

Clone Number: MR3

Alternative Name(s) of Target: SARS CoV 2 S glycoprotein; COVID-19 Spike protein; RBD; Receptor Binding Domain; E2 glycoprotein; E2; Human coronavirus 2 spike glycoprotein; Peplomer protein; S glycoprotein; SARS coronavirus 2 S protein; SARS coronavirus 2 Spike Protein; SARS CoV 2 Spike protein; SARS CoV 2; SARS-CoV-2 S protein; SARSCoV2; SARS-COV-2 S protein; SARS-COV-2 Spike glycoprotein; SARSCOV2 Spike protein; Severe acute respiratory syndrome 2 spike glycoprotein; Severe acute respiratory syndrome virus 2 spike glycoprotein; Spike glycoprotein; 2019-nCoV

UniProt Accession Number of Target Protein: P0DTC2

Published Application(s): biolayer interferometry, FACS, NTRL, therapeutic, ELISA

Published Species Reactivity: SARS Coronavirus 2 (SARS-Cov-2)

Immunogen: The original antibody was selected after one round of ribosome display using three high diversity libraries (concave, loop and convex) followed by three rounds of phage display and panning against biotinylated RBD as a bait and increasing the stringency of selection with every round. The last round of selection was against 5nM RBD.

Specificity: This antibody binds the receptor binding domain (RBD) of the SARS-CoV-2.

Application Notes: MR3 is a potent antibody that binds the RBD with an affinity of $K_D=1.0$ nM and neutralizes SARS-CoV-2 pseudovirus with an IC_{50} of 0.40 µg/mL. Consistent with its highest affinity, MR3 showed the slowest off-rate (2.3×10^{-4} per second). ELISA was used to screen the libraries to identify specific RBD binders. Neutralization activity of the sybody was checked by pre-incubating pseudoviral particles with different concentrations of sybody before infection of VeroE6-hACE2 cells. The rate of infection was then measured by fluorescence-activated cell sorting (FACS). The binding kinetics between sybodies and the RBD was assessed using bio-layer interferometry (Li et al., 2020).

Antibody First Published in: [PMID:](#)

Note on publication:

Product Form

Size:

100 µg Purified antibody.

Purification: Protein A affinity purified

Supplied In: PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -20°C.

Concentration: 1 mg/ml.

Important note – This product is for research use only. It is not intended for use in therapeutic or diagnostic procedures for humans or animals.