

Anti-Monkeypox Virus [VACV-5C7] Standard Size Ab022171.1

Isotype and Format: Mouse IgG1, Kappa

Clone Number: VACV-5C7

Alternative Name(s) of Target: MPXV; MPXV Cph; Monkeypox virus Copenhagen; VACV-5C7 05.11.20;

Anti-ECTV-5C7/1F7; ECTV-5C7/1F7

UniProt Accession Number of Target Protein:

Published Application(s): ELISA, IF

Published Species Reactivity: Monkeypox virus

Immunogen: The original mouse IgG1 version of this antibody was raised against sucrose-gradient purified

monkeypox virus Copenhagen (MPXV Cph).

Specificity: This antibody recognizes monkeypox virus (MPXV). It also crossreacts with other orthopox viruses. Monkeypox virus (MPV) is a double-stranded DNA, zoonotic virus and a species of the genus Orthopoxvirus in the family Poxviridae. It causes a disease that has similar characterists to smallpox but with a milder rash and a lower death rate. The virus causes disease in both humans and animals.

Application Notes: This antibody can be used to identify and analyse Monkypox virus (MPV). This clone was used to identify monkypox virus via ELISA (Czerny et al., 1994; pmid: 7513922). VACV-5C7 can be used to develop diagnostic assays which could allow for the detection of orthopox viruses, especially monkypox virus.

Antibody First Published in: Czerny et al. Epitope detection in the envelope of intracellular naked orthopox viruses and identification of encoding genes Virology. 1994 May 1;200(2):764-77. doi: 10.1006/viro.1994.1240. PMID:7513922

Note on publication: Describes the generation and characterization of antibodies against the Orthopox viruses.

Product Form

Size: 200 μg Purified antibody.

Purification: Protein A affinity purified **Supplied In:** PBS with 0.02% Proclin 300.

Storage Recommendation: Store at 4°C for up to 3 months. For longer storage, aliquot and store at -

20°C.

Concentration:

© 2024 Absolute Antibody	https://absoluteantibody.com/product/anti-monkeypox-virus-vacv-
Important note – This product is for procedures for humans or animals.	research use only. It is not intended for use in therapeutic or diagnostic
1 mg/ml.	